

Technical Data Sheet

Alexa Fluor® 647 Mouse Anti-Stat5 (pY694)


Product Information

Material Number:	612599
Alternate Name:	Signal transducer and activator of transcription 5; MGF; MPF
Size:	50 Tests
Vol. per Test:	20 µl
Clone:	47/Stat5(pY694)
Immunogen:	Phosphorylated Human Phosphorylated Stat5 Peptide
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human Predicted Reactivity: Mouse, Rat, Sheep
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

Stat (Signal transducer and activators of transcription) proteins are critical mediators of the biologic activity of cytokines, including interleukins, interferons, erythropoietin, and growth factors. Ligand-receptor interaction leads to activation of constitutively associated JAK family kinases and subsequent recruitment/activation of Stat proteins by tyrosine phosphorylation. Active Stat proteins then move to the nucleus to promote transcription of cytokine-inducible genes. Seven Stat proteins have been cloned, each of which is differentially expressed and/or activated in a cytokine-specific and cell type-specific manner. Stat5 has been characterized and shown to be encoded by two separate genes, Stat5a and Stat5b that share over 90% identity at the amino acid level. Stat5a has been shown to be involved in lactogenesis and mammary development, while Stat5b has been shown to be involved in growth hormone signaling and to play a role in liver gene expression. Both Stat5a and Stat5b share similarities, both are involved in IL-2 induced peripheral T cell proliferation. The peptide hormone, prolactin, binds to the prolactin receptor (PRLR) to initiate the lactogenic response. There are at least three forms of PRLR; however, only the long form is able to activate the 92-kDa Stat5 protein by inducing phosphorylation at Y694. Once phosphorylated, Stat5 becomes an essential transcription factor which binds to the β-casein gene promoter. The presence of an SH2 domain within Stat5 suggests that it may directly interact with protein tyrosine kinases (PTKs) such as JAK2.

The 47 monoclonal antibody recognizes the phosphorylated Y694 of Stat5a. The homologous phosphorylation site in Stat5b is Y699.



Flow cytometric analysis of Stat5 (pY694). TF-1 cells (Human erythroleukemia; ATCC CRL-2003), a GM-CSF dependent cell line, were starved overnight in IMDM with 0.1% FBS without GM-CSF. The following day, cells were either left unstimulated (unshaded) or stimulated (shaded) with 25 ng/ml GM-CSF recombinant protein (Cat. No. 550068) for 15 minutes at 37°C. Cells were fixed with BD Cytofix™ buffer (10 minutes at 37°C; Cat. No. 554655) and then permeabilized by adding BD Phosflow™ Perm Buffer III (30 minutes on ice; Cat. No. 558050). Cells were then washed twice in BD Pharmingen™ Stain Buffer (Cat. No. 554656), and stained with the Alexa Fluor® 647 Mouse Anti-Stat5 (pY694) antibody. The cells were analyzed on a BD FACSCalibur™ flow cytometer. For intracellular staining of human whole blood, BD Phosflow™ Lyse/Fix buffer (MN 558049) may be used for fixation.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD



Recommended Assay Procedure:

This antibody conjugate is suitable for intracellular staining of human whole blood (using BD Phosflow™ Lyse/Fix Buffer) and peripheral blood mononuclear cells (using BD Cytotfix™ Fixation Buffer or BD Phosflow™ Fix Buffer I).

This purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot analysis (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	PBMC	IL-2	Fixation Buffer	III	Positive Staining
Flow	Human	PBMC	IL-2	Fixation Buffer	I or II	Unsatisfactory
Flow	Human	Whole Blood	IL-2	Lyse/Fix	III	Positive Staining
Flow	Human	Whole Blood	IL-2	Lyse/Fix	I or II	Unsatisfactory
WB	Human	A431 Cell Lysate	EGF	Not Applicable	Not Applicable	92 kDa

Suggested Companion Products

Catalog Number	Name	Size	Clone
550068	Recombinant Human GM-CSF	10 µg	(none)
554655	Fixation Buffer	100 mL	(none)
558050	Perm Buffer III	125 mL	(none)
558049	Lyse/Fix Buffer 5X	250 mL	(none)
557870	Fix Buffer I	250 mL	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100-µl experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.
3. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
4. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
5. Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
7. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
8. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Bromberg J, Darnell JE. The role of STATs in transcriptional control and their impact on cellular function. *Oncogene*. 2000; 19(21):2468-2473. (Biology)

Gouilleux F, Wakao H, Mundt M, Groner B. Prolactin induces phosphorylation of Tyr694 of Stat5 (MGF), a prerequisite for DNA binding and induction of transcription. *EMBO J*. 1994; 13(18):4361-4369. (Biology)

Imada K, Leonard WJ. The Jak-STAT pathway. *Mol Immunol*. 2000; 37:1-11. (Biology)

Johnston RJ, Choi YS, Diamond JA, Yang JA, Crotty S. STAT5 is a potent negative regulator of TFH cell differentiation. *J Exp Med*. 2012; 209(2):243-250. (Clone-specific: Flow cytometry)

Liu KD, Gaffen SL, Goldsmith MA. JAK/STAT signaling by cytokine receptors. *Curr Opin Immunol*. 1998; 10(3):271-278. (Biology)

Prlc M, Bevan MJ. Exploring regulatory mechanisms of CD8+ T cell contraction. *Proc Natl Acad Sci U S A*. 2008; 105(43):16689-16694. (Clone-specific: Flow cytometry)

Suni MA, Maino VC. Flow cytometric analysis of cell signaling proteins. *Methods Mol Biol*. 2011; 717:155-169. (Clone-specific: Flow cytometry)

Wakao H, Gouilleux F, Groner B. Mammary gland factor (MGF) is a novel member of the cytokine regulated transcription factor gene family and confers the prolactin response. *EMBO J*. 1994; 13(9):2182-2191. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

