

Technical Data Sheet

Biotin Mouse Anti-Phosphotyrosine

Product Information


Material Number:	610008
Size:	150 µg
Concentration:	250 µg/ml
Clone:	PY20
Isotype:	Mouse IgG2b
Reactivity:	QC Testing: Human Tested in Development: Chicken, Dog, Frog, Mouse, Rat
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The phosphorylated proteins can be autophosphorylated kinases or certain cellular protein substrates that are regulated in oncogenesis or cell growth. Antibodies to phosphotyrosine provide one of the best tools for the detection and characterization of phosphotyrosine proteins.

Technical Note: The use of milk-containing buffers may interfere with a phosphotyrosine antibody's ability to bind specific proteins of interest. Please use BSA-containing buffers for blocking and incubating purposes.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Western blot analysis of phosphotyrosine on A431 lysate. Lane 1: 1:2000, lane 2: 1:4000, lane 3: 1:8000 dilution of anti-phosphotyrosine, PY20.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at -20° C.

Application Notes

Application

Western blot	Routinely Tested
Immunoprecipitation	Tested During Development
Immunofluorescence	Not Recommended
Immunohistochemistry	Not Recommended

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
611448	A431 + EGF Cell Lysate	500 µg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmlingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Arvidsson AK, Rupp E, Nanberg E, et al. Tyr-716 in the platelet-derived growth factor beta-receptor kinase insert is involved in GRB2 binding and Ras activation. *Mol Cell Biol.* 1994; 14(10):6715-6726.(Biology)

Fan Z, Mendelsohn J, Masui H, Kumar R. Regulation of epidermal growth factor receptor in NIH3T3/HER14 cells by antireceptor monoclonal antibodies. *J Biol Chem.* 1993; 268(28):21073-21079.(Biology)

Glenney JR Jr, Zokas L, Kamps MP. Monoclonal antibodies to phosphotyrosine. *J Immunol Methods.* 1988; 109(2):277-285.(Biology)

Kuppuswamy D, Kerr C, Narishige T, Kasi VS, Menick DR, Cooper G 4th. Association of tyrosine-phosphorylated c-Src with the cytoskeleton of hypertrophying myocardium. *J Biol Chem.* 1997; 272(7):4500-4508.(Clone-specific: Immunoprecipitation)

Nishikawa R, Ji XD, Harmon RC, et al. A mutant epidermal growth factor receptor common in human glioma confers enhanced tumorigenicity. *Proc Natl Acad Sci U S A.* 1994; 91(16):7727-7731.(Biology)