Technical Data Sheet

Purified Mouse Anti-TrkB

Product Information

Material Number: Alternate Name: Size: Concentration: Clone: Immunogen: Isotype: Reactivity: Target MW: Storage Buffer:

610101

Neurotrophic Tyrosine Kinase Receptor Type 2; NTRK2 50 µg 250 µg/ml 47/TrkB Human TrkB aa. 156-322 Mouse IgG1 QC Testing: Rat 116-145 & 70-95 kDa Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

The full-length TrkB gene has been reported to encode for a 145 kDa glycosylated transmembrane tyrosine kinase and neurotrophin receptor. The same gene also has been reported to encode for a 95 kDa glycoprotein that is identical to gp145 [TrkB] at the extracellular domain and transmembrane portion but lacks the intracellular portion. TrkB has been observable in a range of 116-145 kD and 70-95 kD due to various TrkB maturation states, subcellular localizations, and glycosylation states. TrkB belongs to a family of tyrosine kinases that include the TrkA proto-oncogene and TrkC. All have an extracellular ligand-binding domain, a transmembrane region, and intracellular kinase and autophosphorylation domains. TrkB binds the neurotrophins NT3 and NT4/5, as well as brain-derived neurotrophic factor (BDNF), a peptide that helps motor neuron survival and repair. The TrkB tyrosine kinase is activated upon binding to BDNF resulting in autophosphorylation of residues Y670, Y674 and Y675 and the subsequent association of several intracellular proteins like PLC γ , Shc, and PI3-Kinase. TrkB is widely expressed in cells of neuroepithelium and neural crest origin. Some of these include motor neurons, dopamine-producing neurons, and neurons which release γ -aminobutyric acid in the substantia nigra, neocortex, and hippocampus. The two TrkB gene products are differentially expressed in regions of the adult brain.



Western blot analysis of TrkB on a rat cerebrum lysate (left). Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of the Mouse Anti-TrkB antibody.

Preparation and Storage

Store undiluted at -20°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Application Notes

Application						
Western blot	Routinely Tested					
Immunoprecipitation	Not Recommended					
Immunohistochemistry	Not Recommended					
Bioimaging	Not Recommended					
Fluorescence microscopy	Not Recommended					

BD Biosciences

bdbiosciences.c	om				
United States 877.232.8995	Canada 800.268.5430	Europe 32.2.400.98.95	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 55.11.5185.9995
For country cor	tact informatio	on, visit bdbiosci e	ences.com/conta	ct	
Conditions: The in of any patents. BD use of our product product or as a co. written authorizat For Research Use O Unless otherwise r	formation disclose Biosciences will no s. Purchase does n mponent of anoth- tion of Becton, Dic Only. Not for use in noted, BD, BD Logo	d herein is not to be ot be held responsit ot include or carry a er product. Any use kinson and Compan diagnostic or there o and all other trade	e construed as a rec ole for patent infrin ony right to resell o of this product oth y is stictly prohibite peutic procedures. marks are property	ommendation to use gement or other vio r transfer this produc er than the permitte d. Not for resale. • of Becton, Dickinso	e the above product in violation vlations that may occur with the ct either as a stand-alone ad use without the express n and Company. © 2014 BD

Recommended Assav Procedure:

Western blot: Please refer to http://www.bdbiosciences.com/resources/cellbiology/index.jsp

Suggested Companion Products

Catalog Number	Name	Size	Clone
611463	Rat Cerebrum Lysate	500 μg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results. 1
- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 2.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 3.
- discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Armanini MP, McMahon SB, Sutherland J, Shelton DL, Phillips HS. Truncated and catalytic isoforms of trkB are co-expressed in neurons of rat and mouse CNS.. Eur J Neurosci. 1995; 7(6):1403-1409. (Biology)

Du J, Feng L, Yang F, Lu B. Activity- and Ca(2+)-dependent modulation of surface expression of brain-derived neurotrophic factor receptors in hippocampal neurons. J Cell Biol. 2000; 150(6):1423-1434. (Biology: Western blot)

Ebendal T. Function and evolution in the NGF family and its receptors. J Neurosci Res. 1992; 32(4):461-470. (Biology)

Jovanovic JN, Czernik AJ, Fienberg AA, Greengard P, Sihra TS. Synapsins as mediators of BDNF-enhanced neurotransmitter release. Nat Neurosci. 1992; 3(4):323-329. (Biology: Western blot)

Klein R, Conway D, Parada LF, Barbacid M. trkB tyrosine protein kinase gene codes for a second neurogenic receptor that lacks the catalytic kinase domain. Cell. 1990; 61(4):647-656. (Biology)

Kryll D, Yacoubian T, Haapasalo A, Castren E, Lo D, Barker P. Subcellular localization of full-length and truncated trk receptor isoforms in polarized neurons and epithelial cells.. J Neurosci. 1999; 19(14):5823-5833. (Biology)

Muller D, Djebbara-Hannas Z, Jourdain P. Brain-derived neurotrophic factor restores long-term potentiation in polysialic acid-neural cell adhesion

molecule-deficient hippocampus. Proc Natl Acad Sci U S A. 2000; 97(8):4315-4320. (Biology: Western blot)

BD Biosciences

bdbiosciences.com United States

 Canada
 Europe
 Japan

 800.268.5430
 32.2.400.98.95
 0120.8555.90
Asia Pacific 877.232.8995 65.6861.0633 For country contact information, visit bdbiosciences.com/contact

Conditions: The information insclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

Latin America/Caribbean

55.11.5185.999

