Technical Data Sheet

Purified Mouse Anti-Caveolin 3

Product Information

 Material Number:
 610421

 Size:
 150 μg

 Concentration:
 250 μg/ml

 Clone:
 26/Caveolin 3

Immunogen: Rat Caveolin 3 aa. 3-24

 Isotype:
 Mouse IgG1

 Reactivity:
 QC Testing: Rat

Tested in Development: Mouse, Rabbit

Target MW: 18 kDa


Storage Buffer: Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium

azide.


Description

Identified as a tyrosine phosphorylated protein in Rous sarcoma virus-transformed chick embryo fibroblasts (CEF), caveolin is now known to be ubiquitously expressed. Caveolin (also known as VIP21) localizes to non-clathrin membrane invaginations (caveolae) on the inner surface of the plasma membrane. This transmembrane protein plays a structural role in these specializations. Caveolin is also present at the trans-Golgi network (TGN), and similar quantities are found in apically and basolaterally destined transport vesicles. Caveolin is part of a complex containing glycosylphosphatidylinositol (GPI)-linked molecules and cytoplasmic signaling proteins. Caveolin is a transmembrane adaptor molecule that can simultaneously recognize GPI-linked proteins and interact with downstream cytoplasmic signaling molecules, such as c-yes, Annexin II, and hetero-trimeric G proteins. Caveolin 3 has been identified as a distinct isoform which is expressed only in smooth, skeletal, and cardiac muscle.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Western blot analysis of Caveolin 3 on rat muscle. Lane 1: 1:5000, lane 2: 1:10000, lane 3: 1:20000 dilution of anti-Caveolin 3.



Immunofluroescent staining of Rabbit Muscle with anti-Caveolin 3.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20° C.

BD Biosciences

www.bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



610421 Rev. 1 Page 1 of 2

Application Notes

Application

| Western blot | Routinely Tested |
|----------------------|---------------------------|
| Immunohistochemistry | Tested During Development |
| Immunofluorescence | Not Recommended |
| Immunoprecipitation | Not Recommended |

Suggested Companion Products

| Catalog Number | Name | Size | Clone | |
|----------------|--|--------|------------|--|
| 554002 | HRP Goat Anti-Mouse Igs | 1.0 ml | (none) | |
| 554001 | FITC Goat Anti-Mouse Igs (Multiple Adsorption) | 0.5 mg | Polyclonal | |
| 611469 | Rat Muscle Lysate | 500 μg | (none) | |

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Lavitrano M, Bacci ML, Forni M, et al. Efficient production by sperm-mediated gene transfer of human decay accelerating factor (hDAF) transgenic pigs for xenotransplantation. Proc Natl Acad Sci U S A. 2002; 99(22):14230-14235.(Clone-specific: Western blot)

Rybin VO, Xu X, Lisanti MP, Steinberg SF. Differential targeting of beta -adrenergic receptor subtypes and adenylyl cyclase to cardiomyocyte caveolae. A mechanism to functionally regulate the cAMP signaling pathway. J Biol Chem. 2000; 275(52):41447-41457. (Clone-specific: Electron microscopy, Immunoprecipitation, Western blot)

Song KS, Scherer PE, Tang Z, et al. Expression of caveolin-3 in skeletal, cardiac, and smooth muscle cells. Caveolin-3 is a component of the sarcolemma and co-fractionates with dystrophin and dystrophin-associated glycoproteins. J Biol Chem. 1996; 271(25):15160-15165.(Biology)

Woodman SE, Park DS, Cohen AW, et al. Caveolin-3 knock-out mice develop a progressive cardiomyopathy and show hyperactivation of the p42/44 MAPK cascade. J Biol Chem. 2002; 277(41):38988-38997.(Clone-specific: Immunofluorescence, Western blot)

Zhao YY, Liu Y, Stan RV, Fan L, et al. Defects in caveolin-1 cause dilated cardiomyopathy and pulmonary hypertension in knockout mice. Proc Natl Acad Sci U S A. 2002; 99(17):11375-11380.(Clone-specific: Immunohistochemistry, Western blot)

BD Biosciences

www.bdbiosciences.com

United States Canada Asia Pacific Latin America/Caribbean Europe Japan 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



