

## Technical Data Sheet

**Purified Mouse Anti-RIP****Product Information**


|                         |                                                                              |
|-------------------------|------------------------------------------------------------------------------|
| <b>Material Number:</b> | <b>610458</b>                                                                |
| <b>Alternate Name:</b>  | Receptor Interacting Protein                                                 |
| <b>Size:</b>            | 50 µg                                                                        |
| <b>Concentration:</b>   | 250 µg/ml                                                                    |
| <b>Clone:</b>           | 38/RIP                                                                       |
| <b>Immunogen:</b>       | Human RIP aa. 385-650                                                        |
| <b>Isotype:</b>         | Mouse IgG2a                                                                  |
| <b>Reactivity:</b>      | QC Testing: Human<br>Tested in Development: Mouse, Rat, Dog, Chicken         |
| <b>Target MW:</b>       | 74 kDa                                                                       |
| <b>Storage Buffer:</b>  | Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide. |

**Description**

Binding or cross linking of the Fas antigen (also known as APO-1 and CD95) is known to elicit apoptosis in susceptible cells. Fas is a member of a family of cell surface receptors which includes tumor necrosis factor receptors (TNF-R, and TNF-R2) and nerve growth factor receptors (NGF-R), CD40, OX40, CD30, CD27, and 4-1BB. Several members of this family have been shown to regulate or induce cell death (TNF-R1 and TNF-R2). A 74 kDa member of this family protein named RIP (Receptor Interacting Protein) contains an N-terminal region with homology to protein kinases and a C-terminal region containing a cytoplasmic "death domain" present in both Fas and TNF-R1. Both Fas and RIP have been shown to require this death domain to induce apoptosis and overexpression of RIP has been shown to induce cell death in transfected cells.



**Western blot analysis of RIP on a human endothelial cell lysate.** Lane 1: 1:1000, lane 2: 1:2000, lane 3: 1:4000 dilution of the mouse anti-RIP antibody.



**Immunofluorescence staining of WI-38 cells (Human lung fibroblasts; ATCC CCL-75).**

**Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

**BD Biosciences**

bdbiosciences.com

|                      |               |               |              |                     |                                |
|----------------------|---------------|---------------|--------------|---------------------|--------------------------------|
| <b>United States</b> | <b>Canada</b> | <b>Europe</b> | <b>Japan</b> | <b>Asia Pacific</b> | <b>Latin America/Caribbean</b> |
| 877.232.8995         | 888.259.0187  | 32.53.720.550 | 0120.8555.90 | 65.6861.0633        | 55.11.5185.9995                |

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



## Application Notes

### Application

|                      |                           |
|----------------------|---------------------------|
| Western blot         | Routinely Tested          |
| Immunofluorescence   | Tested During Development |
| Immunoprecipitation  | Tested During Development |
| Immunohistochemistry | Not Recommended           |

### Recommended Assay Procedure:

**Western blot:** Please refer to [http://www.bdbiosciences.com/pharmingen/protocols/Western\\_Blotting.shtml](http://www.bdbiosciences.com/pharmingen/protocols/Western_Blotting.shtml)

### Suggested Companion Products

| Catalog Number | Name                          | Size   | Clone      |
|----------------|-------------------------------|--------|------------|
| 611450         | Human Endothelial Cell Lysate | 500 µg | (none)     |
| 554002         | HRP Goat Anti-Mouse Ig        | 1.0 ml | (none)     |
| 554001         | FITC Goat Anti-Mouse Ig       | 0.5 mg | Polyclonal |

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

### References

Devin A, Lin Y, Yamaoka S, Li Z, Karin M, Liu Zg. The alpha and beta subunits of IkappaB kinase (IKK) mediate TRAF2-dependent IKK recruitment to tumor necrosis factor (TNF) receptor 1 in response to TNF. *Mol Cell Biol.* 2002; 21(12):3986-3994.(Biology: Western blot)

Fulda S, Meyer E, Debatin KM. Metabolic inhibitors sensitize for CD95 (APO-1/Fas)-induced apoptosis by down-regulating Fas-associated death domain-like interleukin 1-converting enzyme inhibitory protein expression. *Cancer Res.* 2000; 60(14):3947-3956.(Biology: Western blot)

Lewis J, Devin A, Miller A, et al. Disruption of hsp90 function results in degradation of the death domain kinase, receptor-interacting protein (RIP), and blockage of tumor necrosis factor-induced nuclear factor-kappaB activation. *J Biol Chem.* 2000; 275(14):10519-10526.(Biology: Immunoprecipitation, Western blot)

Stanger BZ, Leder P, Lee TH, Kim E, Seed B. RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell death. *Cell.* 1995; 81(4):513-523.(Biology)

Takahashi T, Tanaka M, Brannan CI, Jenkins NA, Copeland NG, Suda T, and Nagata S. Generalized lymphoproliferative disease in mice, caused by a point mutation in the Fas ligand. *Cell.* 1994; 76(6):969-976.(Biology)