

Anti-Complex I NDUFS3 Monoclonal Antibody

CATALOG #: 439200

COMPONENTS: 100 μg monoclonal antibody

APPLICATIONS: Western blotting

CLONE ID OF MONOCLONAL

ANTIBODY (mAb):

17D950C9H11

SPECIES CROSS-REACTIVITY: human, bovine, rat, mouse, zebrafish, *Drosophila melanogaster*, and *C. elegans*

(+ weaker 60 kDa band)

HOST SPECIES AND ISOTYPE: Mouse IgG2a, κ

IMMUNOGEN: Bovine Heart Complex I

CONCENTRATION: 1 mg/mL in Hepes-Buffered Saline (HBS) with 0.02% azide as a preservative.

SUGGESTED WORKING

CONCENTRATION:

1 μg/mL for Western blotting

mAb PURITY: Near homogeneity as judged by SDS-PAGE. The antibody was produced in

vitro using hybridomas grown in serum-free medium, and then purified by

biochemical fractionation.

STORAGE CONDITIONS: Store at 4°C. Do not freeze.

COUNTRY OF ORIGIN: USA

BACKGROUND:

Complex I, or NADH ubiquinone oxidoreductase, is a large protein complex of 950,000 Da molecular weight made up by 45 to 46 different subunits. A total of seven of the subunits of the complex are encoded by mitochondrial DNA, while the remainder subunits are nuclear encoded, which are translated in the cytosol and translocated into the organelle for assembly at the inner membrane.

The enzyme complex catalyses electron entry from NADH via a flavin (FMN) and several non-heme iron centers. Complex I is sensitive to a wide range of inhibitors, many of which are pesticides or other common environmental toxins, such as rotenone. Complex I dysfunction is a common cause of genetic OXPHOS defects. Altered functioning of this complex is also thought to contribute to several neurological disorders including Parkinson's disease and schizophrenia. Also, there is evidence of Complex I involvement in diabetes.



Related Products

| Product | Conjugate | Cat. No. |
|-----------------------------|--------------|----------|
| Protein A | Sepharose 4B | 10-1041 |
| rec-Protein G | Sepharose 4B | 10-1241 |
| ZyMAX™ Goat anti-rabbit IgG | Unconjugated | 81-6100 |
| ZyMAX™ Goat anti-mouse IgG | Unconjugated | 81-6500 |

Secondary Antibody Conjugates

| Conjugate | Goat anti-rabbit IgG (H+L) | Goat anti-mouse IgG (H+L) | Ex/Em* | Fluorescence similar to |
|------------------|----------------------------|---------------------------|---------|----------------------------|
| Alexa Fluor® 488 | A11008 | A11001 | 495/519 | FITC |
| Alexa Fluor® 555 | A21428 | A21422 | 555/565 | Cy3 |
| Alexa Fluor® 594 | A11012 | A11005 | 590/617 | Texas Red |
| Alexa Fluor® 647 | A21244 | A21235 | 650/668 | Cy5 |
| HRP | 81-6120 | 81-6520 | NA** | NA |
| AP | 81-6122 | 81-6522 | NA | NA |
| Biotin | B2770 | B2763 | NA | NA |

^{*}Excitation/emission (nm); **Not applicable

For additional secondary antibody conjugates, visit www.invitrogen.com/antibodies

For Research Use Only

MitoSciences*

Manufactured exclusively for Invitrogen by MitoSciences, Inc.

 $\underline{www.invitrogen.com}$

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com