



62-8411 1 ml
FITC-Goat Anti-
Human IgG (Gamma)

FITC-Goat Anti-Human IgG (Gamma Chain Specific)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

FORM: Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% BSA, and 0.1% sodium azide.

AFFINITY PURIFIED GOAT ANTI-HUMAN IgG (gamma chain) SPECIFIC ANTIBODY is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and is free of unconjugated FITC.

RECOMMENDED DILUTION: 1:20 to 1:50 dilution with PBS, pH 7.4 containing 10% normal goat serum. Working dilution for a specific application should be determined by individual investigator to obtain the best condition.

APPLICATIONS: Fluorescent antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

STORAGE: Store at 2-8°C.

For research use only.

CAUTION: Not intended for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288
E-mail: techsupport@invitrogen.com

PI628411

(Rev 11/09) DCC-09-1791

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



62-8411 1 ml
FITC-Goat Anti-
Human IgG (Gamma)

FITC-Goat Anti-Human IgG (Gamma Chain Specific)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

FORM: Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% BSA, and 0.1% sodium azide.

AFFINITY PURIFIED GOAT ANTI-HUMAN IgG (gamma chain) SPECIFIC ANTIBODY is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and is free of unconjugated FITC.

RECOMMENDED DILUTION: 1:20 to 1:50 dilution with PBS, pH 7.4 containing 10% normal goat serum. Working dilution for a specific application should be determined by individual investigator to obtain the best condition.

APPLICATIONS: Fluorescent antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

STORAGE: Store at 2-8°C.

For research use only. CAUTION:

Not intended for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288
E-mail: techsupport@invitrogen.com

PI628411

(Rev 11/09) DCC-09-1791

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.