



62-7411 1 ml  
FITC-Goat Anti-  
Human IgA

**FITC-GOAT ANTI-HUMAN IgA (alpha chain specific)**

#### INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

**FORM:** Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% BSA, 40% glycerol and 0.1% sodium azide as preservative.

**AFFINITY PURIFIED GOAT ANTI-HUMAN IgA (alpha chain) SPECIFIC ANTIBODY** is conjugated with fluorescein isothiocyanate (FITC) under optimum condition and is free of unconjugated FITC.

**RECOMMENDED DILUTION:** 1:20 to 1:50 dilution with PBS, pH 7.4 containing 10% normal goat serum. Working dilution for a specific application should be determined by individual investigator to obtain the best condition.

**APPLICATIONS:** Fluorescent-antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

**STORAGE:** Store at 4°C.

**REMARKS:** For research use only. Not for human or drug use. This product has not been assayed for species cross-reactivity unless otherwise indicated

**Research Use Only**

[www.invitrogen.com](http://www.invitrogen.com)

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: [techsupport@invitrogen.com](mailto:techsupport@invitrogen.com)

PI627411

(Rev 10/08) DCC-08-1089

**Important Licensing Information** - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, [www.invitrogen.com](http://www.invitrogen.com)). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



62-7411 1 ml  
FITC-Goat Anti-  
Human IgA

**FITC-GOAT ANTI-HUMAN IgA (alpha chain specific)**

#### INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

**FORM:** Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% BSA, 40% glycerol and 0.1% sodium azide as preservative.

**AFFINITY PURIFIED GOAT ANTI-HUMAN IgA (alpha chain) SPECIFIC ANTIBODY** is conjugated with fluorescein isothiocyanate (FITC) under optimum condition and is free of unconjugated FITC.

**RECOMMENDED DILUTION:** 1:20 to 1:50 dilution with PBS, pH 7.4 containing 10% normal goat serum. Working dilution for a specific application should be determined by individual investigator to obtain the best condition.

**APPLICATIONS:** Fluorescent-antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

**STORAGE:** Store at 4°C.

**REMARKS:** For research use only. Not for human or drug use. This product has not been assayed for species cross-reactivity unless otherwise indicated

**Research Use Only**

[www.invitrogen.com](http://www.invitrogen.com)

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: [techsupport@invitrogen.com](mailto:techsupport@invitrogen.com)

PI627411

(Rev 10/08) DCC-08-1089

**Important Licensing Information** - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, [www.invitrogen.com](http://www.invitrogen.com)). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.