

## **References for Products 21201 to 21204**

1. Boens N, Qin W, Basaric N, Orte A, Talavera EM, Alvarez-Pez JM. (2006) Photophysics of the fluorescent pH indicator BCECF. *J Phys Chem A Mol Spectrosc Kinet Environ Gen Theory*, 110, 9334.
2. Alvarez-Leefmans FJ, Herrera-Perez JJ, Marquez MS, Blanco VM. (2006) Simultaneous measurement of water volume and pH in single cells using BCECF and fluorescence imaging microscopy. *Biophys J*, 90, 608.
3. Bachmeier CJ, Trickler WJ, Miller DW. (2004) Drug efflux transport properties of 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein acetoxyethyl ester (BCECF-AM) and its fluorescent free acid, BCECF. *J Pharm Sci*, 93, 932.
4. Ozkan P, Mutharasan R. (2002) A rapid method for measuring intracellular pH using BCECF-AM. *Biochim Biophys Acta*, 1572, 143.
5. Olson DP, Taylor BJ, Ivy SP. (2001) Detection of MRP functional activity: calcein AM but not BCECF AM as a Multidrug Resistance-related Protein (MRP1) substrate. *Cytometry*, 46, 105.
6. Marechal X, Mordon S, Devoisselle JM, Begu S, Guery B, Neviere R, Buys B, Dhelin G, Lesage JC, Mathieu D, Chopin C. (1999) In vivo application of intestinal pH measurement using 2',7'-bis(carboxyethyl)-5,6-carboxyfluorescein (BCECF) fluorescence imaging. *Photochem Photobiol*, 70, 813.
7. Ruiz-Palomo F, Garcia C, Gomez M, Revenga M. (1999) Fluorescence probe (BCECF) loading efficiency in human platelets depends on cell concentration: application to pH<sub>i</sub> measurements. *Clin Biochem*, 32, 391.
8. Weinlich M, Heydasch U, Mooren F, Starlinger M. (1998) Simultaneous detection of cell volume and intracellular pH in isolated rat duodenal cells by confocal microscopy and BCECF. *Res Exp Med (Berl)*, 198, 73.
9. Scott DA, Docampo R, Benchimol M. (1998) Analysis of the uptake of the fluorescent marker 2',7'-bis-(2-carboxyethyl)-5-and-6-carboxyfluorescein (BCECF) by hydrogenosomes in *Trichomonas vaginalis*. *Eur J Cell Biol*, 76, 139.
10. Sandhu V, Miller M, Grover AK. (1998) Effects of peroxide on the fluorescence of the Ca<sup>2+</sup> probe Fluo 3 and the pH probe BCECF. *Mol Cell Biochem*, 178, 77.
11. Weinlich M, Theiss C, Lin CT, Kinne RK. (1998) BCECF in single cultured cells: inhomogeneous distribution but homogeneous response. *J Exp Biol*, 201, 57.
12. Grant RL, Acosta D. (1997) Ratiometric measurement of intracellular pH of cultured cells with BCECF in a fluorescence multi-well plate reader. *In Vitro Cell Dev Biol Anim*, 33, 256.
13. Devoisselle JM, Soulie S, Mordon S, Maillols H. (1996) Fluorescent characteristics and pharmacokinetic profiles of the fluorescent probe BCECF in various tissues: the role of blood content. *Photochem Photobiol*, 64, 906.
14. Aharonovitz O, Fridman H, Livne AA, Granot Y. (1996) The effect of BCECF on intracellular pH of human platelets. *Biochim Biophys Acta*, 1284, 227.
15. Boyarsky G, Hanssen C, Clyne LA. (1996) Inadequacy of high K<sup>+</sup>/nigericin for calibrating BCECF. II. Intracellular pH dependence of the correction. *Am J Physiol*, 271, C1146.
16. Boyarsky G, Hanssen C, Clyne LA. (1996) Inadequacy of high K<sup>+</sup>/nigericin for calibrating BCECF. I. Estimating steady-state intracellular pH. *Am J Physiol*, 271, C1131.
17. Boyarsky G, Hanssen C, Clyne LA. (1996) Superiority of in vitro over in vivo calibrations of BCECF in vascular smooth muscle cells. *Faseb J*, 10, 1205.
18. Franck P, Petitipain N, Cherlet M, Dardennes M, Maachi F, Schutz B, Poisson L, Nabot P. (1996) Measurement of intracellular pH in cultured cells by flow cytometry with BCECF-AM. *J Biotechnol*, 46, 187.
19. Bevensee MO, Schwiening CJ, Boron WF. (1995) Use of BCECF and propidium iodide to assess membrane integrity of acutely isolated CA1 neurons from rat hippocampus. *J Neurosci Methods*, 58, 61.

20. Obexer W, Schmid C, Brun R. (1995) A novel in vitro screening assay for trypanocidal activity using the fluorescent dye BCECF-AM. *Trop Med Parasitol*, 46, 45.
21. Carpenter L, Halestrap AP. (1994) The kinetics, substrate and inhibitor specificity of the lactate transporter of Ehrlich-Lettre tumour cells studied with the intracellular pH indicator BCECF. *Biochem J*, 304 ( Pt 3), 751.
22. Wang X, Levi AJ, Halestrap AP. (1994) Kinetics of the sarcolemmal lactate carrier in single heart cells using BCECF to measure pH<sub>i</sub>. *Am J Physiol*, 267, H1759.
23. King GG, Lohrmann WE, Ickes JW, Jr., Feldman GM. (1994) Identification of Na<sup>+</sup>/H<sup>+</sup> exchange on the apical side of surface colonocytes using BCECF. *Am J Physiol*, 267, G119.
24. Futsaether CM, Kjeldstad B, Johnsson A. (1993) Measurement of the intracellular pH of Propionibacterium acnes: comparison between the fluorescent probe BCECF and 31P-NMR spectroscopy. *Can J Microbiol*, 39, 180.
25. James-Krake MR. (1992) Quick and accurate method to convert BCECF fluorescence to pH<sub>i</sub>: calibration in three different types of cell preparations. *J Cell Physiol*, 151, 596.
26. Allen CN, Harpur ES, Gray TJ, Simmons NL, Hirst BH. (1990) Efflux of bis-carboxyethyl-carboxyfluorescein (BCECF) by a novel ATP-dependent transport mechanism in epithelial cells. *Biochem Biophys Res Commun*, 172, 262.
27. Borzak S, Kelly RA, Kramer BK, Matoba Y, Marsh JD, Reers M. (1990) In situ calibration of fura-2 and BCECF fluorescence in adult rat ventricular myocytes. *Am J Physiol*, 259, H973.
28. Mozingo NM, Chandler DE. (1990) The fluorescent probe BCECF has a heterogeneous distribution in sea urchin eggs. *Cell Biol Int Rep*, 14, 689.
29. Noel J, Tejedor A, Vinay P, Laprade R. (1989) Fluorescence measurement of intracellular pH on proximal tubule suspensions. The need for a BCECF sink. *Ren Physiol Biochem*, 12, 371.
30. Miyata H, Hayashi H, Suzuki S, Noda N, Kobayashi A, Fujiwake H, Hirano M, Yamazaki N. (1989) Dual loading of the fluorescent indicator fura-2 and 2,7-biscarboxyethyl-5(6)-carboxyfluorescein (BCECF) in isolated myocytes. *Biochem Biophys Res Commun*, 163, 500.
31. Kapus A, Ligeti E, Fonyo A. (1989) Na<sup>+</sup>/H<sup>+</sup> exchange in mitochondria as monitored by BCECF fluorescence. *FEBS Lett*, 251, 49.
32. Kawase T, Suzuki A. (1989) Fluoride-induced cytoplasmic acidification: possible role of protein kinase C in BCECF-loaded L929 cells. *Pharmacol Toxicol*, 64, 426.
33. Weiner ID, Hamm LL. (1989) Use of fluorescent dye BCECF to measure intracellular pH in cortical collecting tubule. *Am J Physiol*, 256, F957.
34. Siffert W, Siffert G, Scheid P, Akkerman JW. (1989) Activation of Na<sup>+</sup>/H<sup>+</sup> exchange and Ca<sup>2+</sup> mobilization start simultaneously in thrombin-stimulated platelets. Evidence that platelet shape change disturbs early rises of BCECF fluorescence which causes an underestimation of actual cytosolic alkalinization. *Biochem J*, 258, 521.
35. Kolber MA, Quinones RR, Gress RE, Henkart PA. (1988) Measurement of cytotoxicity by target cell release and retention of the fluorescent dye bis-carboxyethyl-carboxyfluorescein (BCECF). *J Immunol Methods*, 108, 255.
36. Davis MH, Altschuld RA, Jung DW, Brierley GP. (1987) Estimation of intramitochondrial pCa and pH by fura-2 and 2,7 biscarboxyethyl-5(6)-carboxyfluorescein (BCECF) fluorescence. *Biochem Biophys Res Commun*, 149, 40.