References for Product 21495

- 1. Srinivasan G, Friauf E, Lohrke S. (2004) Functional glutamatergic and glycinergic inputs to several superior olivary nuclei of the rat revealed by optical imaging. Neuroscience, 128, 617.
- 2. Obaid AL, Loew LM, Wuskell JP, Salzberg BM. (2004) Novel naphthylstyryl-pyridium potentiometric dyes offer advantages for neural network analysis. J Neurosci Methods, 134, 179.
- 3. Fisher JA, Civillico EF, Contreras D, Yodh AG. (2004) In vivo fluorescence microscopy of neuronal activity in three dimensions by use of voltage-sensitive dyes. Opt Lett, 29, 71.
- 4. Straub H, Kuhnt U, Hohling JM, Kohling R, Gorji A, Kuhlmann D, Tuxhorn I, Ebner A, Wolf P, Pannek HW, Lahl R, Speckmann EJ. (2003) Stimulus-induced patterns of bioelectric activity in human neocortical tissue recorded by a voltage sensitive dye. Neuroscience, 121, 587.
- 5. Senner V, Kohling R, Puttmann-Cyrus S, Straub H, Paulus W, Speckmann EJ. (2004) A new neurophysiological/neuropathological ex vivo model localizes the origin of gliomaassociated epileptogenesis in the invasion area. Acta Neuropathol (Berl), 107, 1.
- 6. Hulsmann S, Straub H, Richter DW, Speckmann EJ. (2003) Blockade of astrocyte metabolism causes delayed excitation as revealed by voltage-sensitive dyes in mouse brainstem slices. Exp Brain Res, 150, 117.
- 7. Horikawa J, Hess A, Nasu M, Hosokawa Y, Scheich H, Taniguchi I. (2001) Optical imaging of neural activity in multiple auditory cortical fields of guinea pigs. Neuroreport, 12, 3335.
- 8. Takuma S. (2001) Effect of neonatal capsaicin treatment on neural activity in the medullary dorsal horn of neonatal rats evoked by electrical stimulation to the trigeminal afferents: an optical, electrophysiological, and quantitative study. Brain Res, 906, 1.
- 9. Le Calvez S, Ulfendahl M. (2000) An in vitro preparation to access cellular and neuronal components in the mouse inner ear. J Neurocytol, 29, 645.
- 10. Maeda S, Inagaki S, Kawaguchi H, Song WJ. (2001) Separation of signal and noise from in vivo optical recording in Guinea pigs using independent component analysis. Neurosci Lett, 302, 137.
- 11. Flock A, Flock B, Scarfone E. (1998) Laser scanning confocal microscopy of the hearing organ: fluorochrome-dependent cellular damage is seen after overexposure. J Neurocytol, 27, 507.
- 12. Cohen D, Yarom Y. (1999) Optical measurements of synchronized activity in isolated mammalian cerebellum. Neuroscience, 94, 859.
- 13. de Curtis M, Takashima I, lijima T. (1999) Optical recording of cortical activity after in vitro perfusion of cerebral arteries with a voltage-sensitive dye. Brain Res, 837, 314.
- 14. Imaizumi M, Oguma Y, Kawatani M. (1998) Optical imaging of the spontaneous neuronal activities in the male rat major pelvic ganglion following denervation of the pelvic nerve. Neurosci Lett, 258, 159.
- 15. Galizia CG, Nagler K, Holldobler B, Menzel R. (1998) Odour coding is bilaterally symmetrical in the antennal lobes of honeybees (Apis mellifera). Eur J Neurosci, 10, 2964.
- 16. Moriyama T, Horikawa J, Hosokawa Y, Taniguchi I. (1997) Optical recording of azimuth representation in guinea pig auditory cortex. Acta Otolaryngol Suppl, 532, 138.
- 17. Kleinfeld D, Delaney KR. (1996) Distributed representation of vibrissa movement in the upper layers of somatosensory cortex revealed with voltage-sensitive dyes. J Comp Neurol, 375, 89.
- 18. Onimaru H, Kanamaru A, Homma I. (1996) Optical imaging of respiratory burst activity in newborn rat medullary block preparations. Neurosci Res, 25, 183.
- 19. Langenstroth M, Albowitz B, Kuhnt U. (1996) Partial suppression of GABAA-mediated inhibition induces spatially restricted epileptiform activity in guinea pig neocortical slices. Neurosci Lett, 210, 103.

20. Elias SA, Yae H, Ebner TJ. (1993) Optical imaging of parallel fiber activation in the rat cerebellar cortex: spatial effects of excitatory amino acids. Neuroscience, 52, 771.