

## References for Product 5524

1. Kipper MJ, Kleinman HK, Wang FW. (2007) Covalent surface chemistry gradients for presenting bioactive peptides. *Anal Biochem*, 363, 175.
2. Gan J, Harper TW, Hsueh MM, Qu Q, Humphreys WG. (2005) Dansyl glutathione as a trapping agent for the quantitative estimation and identification of reactive metabolites. *Chem Res Toxicol*, 18, 896.
3. Christiansen C, St Hilaire PM, Winther JR. (2004) Fluorometric polyethyleneglycol-peptide hybrid substrates for quantitative assay of protein disulfide isomerase. *Anal Biochem*, 333, 148.
4. Storhoff JJ, Marla SS, Bao P, Hagenow S, Mehta H, Lucas A, Garimella V, Patno T, Buckingham W, Cork W, Muller UR. (2004) Gold nanoparticle-based detection of genomic DNA targets on microarrays using a novel optical detection system. *Biosens Bioelectron*, 19, 875.
5. Bayle C, Issac C, Salvayre R, Couderc F, Causse E. (2002) Assay of total homocysteine and other thiols by capillary electrophoresis and laser-induced fluorescence detection. II. Pre-analytical and analytical conditions. *J Chromatogr A*, 979, 255.
6. Raje S, Glynn NM, Thorpe C. (2002) A continuous fluorescence assay for sulfhydryl oxidase. *Anal Biochem*, 307, 266.
7. Causse E, Issac C, Malatray P, Bayle C, Valdiguié P, Salvayre R, Couderc F. (2000) Assays for total homocysteine and other thiols by capillary electrophoresis-laser-induced fluorescence detection. I. Preanalytical condition studies. *J Chromatogr A*, 895, 173.
8. Causse E, Malatray P, Calaf R, Charpiot P, Candito M, Bayle C, Valdiguié P, Salvayre R, Couderc F. (2000) Plasma total homocysteine and other thiols analyzed by capillary electrophoresis/laser-induced fluorescence detection: comparison with two other methods. *Electrophoresis*, 21, 2074.
9. Salazar JF, Schorr H, Herrmann W, Herbeth B, Siest G, Leroy P. (1999) Measurement of thiols in human plasma using liquid chromatography with precolumn derivatization and fluorescence detection. *J Chromatogr Sci*, 37, 469.
10. Causse E, Terrier R, Champagne S, Nertz M, Valdiguié P, Salvayre R, Couderc F. (1998) Quantitation of homocysteine in human plasma by capillary electrophoresis and laser-induced fluorescence detection. *J Chromatogr A*, 817, 181.
11. Tyurin VA, Tyurina YY, Quinn PJ, Schor NF, Balachandran R, Day BW, Kagan VE. (1998) Glutamate-induced cytotoxicity in PC12 pheochromocytoma cells: role of oxidation of phospholipids, glutathione and protein sulfhydryls revealed by bcl-2 transfection. *Brain Res Mol Brain Res*, 60, 270.
12. Kok RJ, Visser J, Moolenaar F, de Zeeuw D, Meijer DK. (1997) Bioanalysis of captopril: two sensitive high-performance liquid chromatographic methods with pre- or postcolumn fluorescent labeling. *J Chromatogr B Biomed Sci Appl*, 693, 181.
13. Bellomo G, Palladini G, Vairetti M. (1997) Intranuclear distribution, function and fate of glutathione and glutathione-S-conjugate in living rat hepatocytes studied by fluorescence microscopy. *Microsc Res Tech*, 36, 243.
14. Ercal N, Oztezcan S, Hammond TC, Matthews RH, Spitz DR. (1996) High-performance liquid chromatography assay for N-acetylcysteine in biological samples following derivatization with N-(1-pyrenyl)maleimide. *J Chromatogr B Biomed Appl*, 685, 329.
15. Daskalakis I, Luccock MD, Anderson A, Wild J, Schorah CJ, Levene MI. (1996) Determination of plasma total homocysteine and cysteine using HPLC with fluorescence detection and an ammonium 7-fluoro-2, 1, 3-benzoxadiazole-4-sulphonate (SBD-F) derivatization protocol optimized for antioxidant concentration, derivatization reagent concentration, temperature and matrix pH. *Biomed Chromatogr*, 10, 205.
16. Ji AJ, Savon SR, Jacobsen DW. (1995) Determination of total serum sulfite by HPLC with fluorescence detection. *Clin Chem*, 41, 897.
17. Fadden P, Haystead TA. (1995) Quantitative and selective fluorophore labeling of phosphoserine on peptides and proteins: characterization at the attomole level by capillary electrophoresis and laser-induced fluorescence. *Anal Biochem*, 225, 81.

18. Rabel SR, Stobaugh JF, Trueworthy R. (1995) Determination of intracellular levels of 6-mercaptapurine metabolites in erythrocytes utilizing capillary electrophoresis with laser-induced fluorescence detection. *Anal Biochem*, 224, 315.
19. O'Keefe DO. (1994) Quantitative electrophoretic analysis of proteins labeled with monobromobimane. *Anal Biochem*, 222, 86.
20. Bolton RM, Haritos VS, Whitehouse MW, Ahokas JT. (1994) Ammonium 4-chloro-7-sulfobenzofurazan: a fluorescent substrate highly specific for rat glutathione S-transferase subunit 3. *Anal Biochem*, 216, 418.
21. Scott RB, Collins JM, Matin S, White F, Swerdlow PS. (1990) Simultaneous measurement of neutrophil, lymphocyte, and monocyte glutathione by flow cytometry. *J Clin Lab Anal*, 4, 324.
22. Jacobsen DW, Gatautis VJ, Green R. (1989) Determination of plasma homocysteine by high-performance liquid chromatography with fluorescence detection. *Anal Biochem*, 178, 208.
23. Cotgreave IA, Weis M, Berggren M, Sandy MS, Moldeus PW. (1988) Determination of the intracellular protein thiol distribution of hepatocytes using monobromobimane derivatisation of intact cells and isolated subcellular fractions. *J Biochem Biophys Methods*, 16, 247.
24. Cotgreave IA, Moldeus P. (1986) Methodologies for the application of monobromobimane to the simultaneous analysis of soluble and protein thiol components of biological systems. *J Biochem Biophys Methods*, 13, 231.
25. Ayers FC, Warner GL, Smith KL, Lawrence DA. (1986) Fluorometric quantitation of cellular and nonprotein thiols. *Anal Biochem*, 154, 186.
26. Broekaert D, Coucke P, Nsabumukunzi S, Reyniers P, van Oostveldt P, Kluyskens P, Gillis E. (1982) Quantitative determination of free thiol and disulphide groups by a fluorescent maleimide procedure. *Acta Histochem*, 70, 62.
27. Sippel TO. (1981) Microfluorometric analysis of protein thiol groups with a coumarinylphenylmaleimide. *J Histochem Cytochem*, 29, 1377.